

Comparison of superconductivity in Sr_2RuO_4 and copper oxides

E.V.Kuz'min* and S.G.Ovchinnikov**

* Krasnoyarsk State University, Svobodnyi av. 74, Krasnoyarsk 660062, Russia

** L.V.Kirensky Institute of Physics, Krasnoyarsk, 660036, Russia

(February 1, 2008)

To compare the superconductivity in strongly correlated electron systems with the antiferromagnetic fluctuations in the copper oxides and with the ferromagnetic fluctuations in Sr_2RuO_4 a $t-J-I$ model is proposed. The antiferromagnetic coupling J results in the superconducting state of $d_{x^2-y^2}$ symmetry and the ferromagnetic coupling I results in the spin-triplet p -type state. The difference in the gap anisotropies provides the large difference in T_c values, for the typical values of the coupling constants: $T_c \sim 1K$ for the ruthenate and $T_c \sim 100K$ for the cuprates.

PACS numbers: 71.27.+a, 74.25

More than a decade of intensive research of the cuprate superconductors and related systems has raised fundamental challenges to our understanding of the mechanism of high-temperature superconductivity (SC). One of the most important question is what is so specific in copper oxides, is it the unique chemistry of the planar $Cu-O$ bond that determines the high value of T_c ? The discovery of SC in Sr_2RuO_4 with $T_c \sim 1K$ [1] is of a particular interest because it has a similar crystal structure to the parent compound La_2CuO_4 , of one of the best studied families of the cuprate superconductors, $La_{2-x}Sr_xCuO_4$, but has four valence electrons (for Ru^{4+}) instead of one hole per formula unit. It is generally believed that comparison of normal and SC properties of the cuprates and the ruthenate will give more deeper understanding of the nature of high- T_c SC. While the normal state of doped cuprates looks like almost antiferromagnetic Fermi-liquid [2], the normal state of Sr_2RuO_4 is characterised by the strong ferromagnetic fluctuations [3]. Properties of SC state are also different: the singlet pairing with major contribution of the $d_{x^2-y^2}$ symmetry was suggested for the cuprates [4], while the triplet pairing with p -type symmetry similar to the 3He A_1 phase is proposed for Sr_2RuO_4 [5]. The triplet SC in Sr_2RuO_4 is induced by the ferromagnetic spin fluctuations [6].

To compare the SC in Sr_2RuO_4 and cuprates we have proposed here a $t-J-I$ model containing both an indirect antiferromagnetic coupling J and a direct ferromagnetic coupling I between neighboring cations. This model is based on the electronic structure calculations. An

important difference from the cuprates is that relevant orbitals to the states near the Fermi energy are Ru $d\epsilon(d_{xy}, d_{yz}, d_{xz})$ and O $p\pi$, instead of Cu $d_{x^2-y^2}$ and O $p\sigma$ states. Due to σ -bonding in the cuprates a strong $p-d$ hybridization takes place resulting in the strong antiferromagnetic coupling J , a direct $d_{x^2-y^2}$ $Cu-Cu$ overlapping is negligible. In Sr_2RuO_4 with π -bonding the $Ru-O-Ru$ 180-degree antiferromagnetic superexchange coupling is weak [7] while a direct d_{xy} $Ru-Ru$

overlapping is not small. That is why we add the Heisenberg type direct $Ru-Ru$ exchange interaction to the Hamiltonian of the $t-J$ model. The strong electron correlations are common features of the charge carriers both in cuprates and Sr_2RuO_4 in our model. These correlations for the cuprates are well known [8]. The importance of electron correlations for Sr_2RuO_4 follows from the high value of the effective mass of electrons in the γ -band obtained by the quantum oscillations measurements [9].

We have found the different solutions for SC state in $t-J-I$ model: with the singlet d -type pairing governed by the antiferromagnetic coupling J and with the triplet p -type pairing induced by the ferromagnetic coupling I . The equations for T_c in both states are similar. Nevertheless the same absolute value of the coupling constants results in quite different T_c values, $T_c^{(p)} \sim 1K$ and $T_c^{(d)} \sim 100K$ for typical values of parameters. The gap anisotropy is responsible for the large difference in the T_c values. For the p -type pairing the $\mathbf{k}$ -dependence of the gap provides cancellation of the singular van-Hove contribution of the two-dimensional density of states, while the $\mathbf{k}$ -dependence of the d -type gap results in the significant contribution of the van-Hove singularity.

The Hamiltonian of the $t-J-I$ model is written in the form

$$H = \sum_{\mathbf{f}\sigma} (\varepsilon - \mu) X_{\mathbf{f}}^{\sigma\sigma} - t \sum_{\mathbf{f}\delta\sigma} X_{\mathbf{f}}^{\sigma 0} X_{\mathbf{f}+\delta}^{0\sigma} + J \sum_{\mathbf{f}\delta} K_{\mathbf{f},\mathbf{f}+\delta}^{(-)} - I \sum_{\mathbf{f}\delta} K_{\mathbf{f},\mathbf{f}+\delta}^{(+)} , \quad (1)$$

$$K_{\mathbf{f}\mathbf{g}}^{(\pm)} = \vec{S}_{\mathbf{f}} \cdot \vec{S}_{\mathbf{g}} \pm \frac{1}{4} n_{\mathbf{f}} n_{\mathbf{g}} , \quad X_{\mathbf{f}}^{\uparrow\uparrow} + X_{\mathbf{f}}^{\downarrow\downarrow} + X_{\mathbf{f}}^{00} = 1.$$

Here the Hubbard X-operators $X_{\mathbf{f}}^{pq} = |p\rangle\langle q|$ are determined in the reduced Hilbert space containing empty states $|0\rangle$ and single-occupied states $|\sigma\rangle$, $\sigma = \uparrow$ or $\sigma = \downarrow$. The X-operators algebra exactly takes into accounts the




FIG. 1. Concentration dependence of T_c for the triplet pairing of the p -type.

constrain condition that is one of the important effects of the strong electron correlations. The $\vec{S}_f$ and n_f operators in (1) are the usual spin and number of particles operators at the site f , δ is the vector between n.n.. For the cuprates $J \gg I$, but for Sr_2RuO_4 $I \gg J$.

To get SC the copper oxides should be doped while Sr_2RuO_4 is self-doped. According to the band structure calculations [10] the electron α -band in Sr_2RuO_4 is half-filled, the hole β -band has $n_0 = 0.28$ holes and the electron γ -band with d_{xy} contribution is more then half-filled, $n_\gamma = 1 + n_0$. The strong electron correlations split the γ -band into filled lower Hubbard band (LHB) with $n_e = 1$ and partially filled upper Hubbard band (UHB) with the electron concentration $n_e = n_0$. We use the hole representation where the electron UHB transforms in the hole LHB with hole concentration $n_h = 1 - n_0$. All other bands (α and β) are treated here as an electron reservoir. Observation of a square flux-line lattice in Sr_2RuO_4 allows to suggest that SC resides mainly on the γ band [11]. For the cuprates the quasiparticle is a hole in the electron LHB with the electron concentration $n_e = 1 - n_0$, for $La_{2-x}Sr_xCuO_4$ $n_0 = x$. There are many ways to get the mean-field solutions for SC state, we have used the irreducible Green function method [12] projecting the higher-order Green functions onto subspace of normal $\langle\langle X_k^{0\sigma} | X_k^{\sigma 0} \rangle\rangle$ and abnormal $\langle\langle X_k^{-\sigma 0} X_k^{\sigma 0} \rangle\rangle$ Green functions coupled via the Gorkov system of equations. Three different solutions have been studied: singlet s - and d - and triplet p -types. The gap equation has the form for s -state

$$1 = \frac{1}{N} \sum_{\mathbf{p}} \frac{2\omega_{\mathbf{p}} + (2g - \lambda)\omega_{\mathbf{p}}^2}{2E_{\mathbf{p}0}} \tanh\left(\frac{E_{\mathbf{p}0}}{2\tau}\right) \quad (2)$$

and for the p, d -states




FIG. 2. Concentration dependence of T_c for the singlet pairing of the d -type.

$$\frac{1}{\alpha_l} = \frac{1}{N} \sum_{\mathbf{p}} \frac{\psi_l^2(\mathbf{p})}{2E_{\mathbf{p}l}} \tanh\left(\frac{E_{\mathbf{p}l}}{2\tau}\right). \quad (3)$$

Here for s, p, d -states

$$E_{\mathbf{p}l} = \sqrt{c^2(n_0)(\omega_{\mathbf{p}} - m)^2 + |\Delta_{\mathbf{p}l}|^2},$$

$$m = \left[\frac{\mu - \epsilon}{zt} + (g + \lambda) \frac{1 - n_0}{2} \right] / c(n_0),$$

where $\omega_{\mathbf{p}} = -\gamma_{\mathbf{p}} = -(1/z) \sum_{\delta} \exp(i\mathbf{p}\delta)$, m is a dimensionless chemical potential, $c(n_0) = (1 + n_0)/2$, $g = J/t$, $\lambda = I/t$ and $\tau = k_B T / zt$ is a dimensionless temperature. The SC gaps are equal to

$$\Delta_{\mathbf{k}0} = [2 + (2g - \lambda)\omega_{\mathbf{k}}] \Delta_0, \Delta_0 = \frac{1}{N} \sum_{\mathbf{p}} \omega_{\mathbf{p}} B_{\mathbf{p}} / c(n_0),$$

$$B_{\mathbf{p}} = \langle X_{-\mathbf{p}}^{0l} X_{\mathbf{p}}^{0l} \rangle \quad (4)$$

for s -type ($l = 0$) and

$$\Delta_{\mathbf{k}l} = \alpha_l \psi_l(\mathbf{k}) \frac{1}{N} \sum_{\mathbf{p}} \psi_l(\mathbf{p}) B_{\mathbf{p}} / c(n_0) \quad (5)$$

for the p - and d - states. The coupling constants and the gap anisotropy in the l -th channel are given by

$$\alpha_p = \lambda, \quad \psi_p(\mathbf{k}) = \frac{1}{2}(\sin k_x + \sin k_y), \quad (6)$$

$$\alpha_d = (2g - \lambda), \quad \psi_d(\mathbf{k}) = \frac{1}{2}(\cos k_x - \cos k_y). \quad (7)$$

Here we have considered only the two-dimensional square lattice with the lattice parameter $a = 1$. The equation for the chemical potential has the form

$$1 - n_0 = \frac{1}{N} \sum_{\mathbf{k}\sigma} \langle X_{\mathbf{k}}^{\sigma 0} X_{\mathbf{k}}^{0\sigma} \rangle. \quad (8)$$

An important effect of the strong electron correlation is the constrain condition excluding double-occupied states

$$\frac{1}{N} \sum_{\mathbf{k}} B_{\mathbf{k}} = \frac{1}{N} \sum_{\mathbf{k}} \langle X_{-\mathbf{k}}^{0\downarrow} X_{\mathbf{k}}^{0\uparrow} \rangle = 0. \quad (9)$$

The first term in (2) for the singlet s -type pairing parameter is proportional to $2tz$ and appears due to the kinematic mechanism of pairing [13]. The s -type solution does not satisfy to the constrain condition (9) [14] while for p - and d -type it is fulfilled. The equation for T_c in p - and d -states is given by

$$\frac{2c(n_0)}{\alpha_l} = \frac{1}{N} \sum_{\mathbf{p}} \frac{\psi_l^2(\mathbf{p})}{|\omega_{\mathbf{p}} - m|} \tanh \left(\frac{c(n_0)|\omega_{\mathbf{p}} - m|}{2\tau_c^{(l)}} \right). \quad (10)$$

The same equation for the $d_{x^2-y^2}$ -pairing has been derived by the diagram technique for the $t - J$ model [15].

At the numerical solution of the equations (10) more than 10^6 points of the Brillouine zone have been taken. Results of $T_c(n_0)$ computations are shown in the Fig.1 and 2 for several values of the coupling constants α_l . These results have revealed the remarkable difference in T_c values: $T_c^{(p)} \ll T_c^{(d)}$ when $\alpha_p = \alpha_d$. The moderate values of $\alpha \approx 0.4 - 0.5$ and $zt \approx 0.5eV$ result in $T_c^{(p)} \sim 1K, T_c^{(d)} \sim 100K$.

It is clear from equations (6),(7) that the p -type SC is formed by the ferromagnetic interaction, that is the case of Sr_2RuO_4 , and the d -type SC is induced by the antiferromagnetic interaction in copper oxides. To understand why $T_c^{(d)} \gg T_c^{(p)}$ we have analysed the eqn. (10) analytically. Using integration over the constant energy surfaces $\omega_{\mathbf{k}} = \omega$ it can be rewritten like

$$\frac{2c(n_0)}{\alpha_l} = \int_{-1}^{+1} \frac{\psi_l^2(\omega)}{|\omega - m|} \tanh \left(\frac{c(n)|\omega - m|}{2\tau_c} \right) d\omega, \quad (11)$$

$$\psi_l^2(\omega) = \frac{1}{(2\pi)^2} \oint_{(\sigma_{\omega})} \frac{\psi_l^2(\mathbf{k})}{|\nabla_{\mathbf{k}}\omega_{\mathbf{k}}|} d\sigma_{\omega}. \quad (12)$$

The sum rule for the $\psi_l^2(\omega)$ functions is the same for $l = p$ and $l = d$:

$$\frac{1}{N} \sum_{\mathbf{k}} \psi_l^2(\mathbf{k}) = \int_{-1}^{+1} \psi_l^2(\omega) d\omega = 1/4. \quad (13)$$

For the p -state

$$\frac{\psi_p^2(\mathbf{k})}{|\nabla_{\mathbf{k}}\omega_{\mathbf{k}}|} = |\nabla_{\mathbf{k}}\omega_{\mathbf{k}}| + \frac{\sin k_x \sin k_y}{\sqrt{\sin^2 k_x + \sin^2 k_y}} \quad (14)$$

the second term in (14) gives zero contribution to the integral (12) and $\psi_p^2(\omega)$ is rather small with smooth energy dependence, $\psi_p^2(\omega) \approx (2/\pi^2)(1 - |\omega|^{1.61})$. For the d -state

$$\frac{\psi_d^2(\mathbf{k})}{|\nabla_{\mathbf{k}}\omega_{\mathbf{k}}|} = \frac{1}{2} \frac{(\cos k_x - \cos k_y)^2}{\sqrt{\sin^2 k_x + \sin^2 k_y}} \quad (15)$$

has the same singularity as the van-Hove singularity in the density of states $\rho(\omega)$. The result of calculation is

$$\psi_d^2(\omega) = (1 - \omega^2)\rho(\omega) - 2\psi_p^2(\omega),$$

$$\rho(\omega) = \frac{1}{\pi} - \left(\frac{1}{2} - \frac{1}{\pi} \right) \ln(|\omega|). \quad (16)$$

The comparison of $\psi_p^2(\omega)$ and $\psi_d^2(\omega)$ has shown that the van-Hove singularity is cancelled in the p -state and does not cancelled in the d -state (Fig 3).




FIG. 3. Energy dependence of the effective gap anisotropy ψ_p^2 and ψ_d^2 and single-particle density of states $\rho(\omega)$ in square lattice.

In conclusion we have presented the model of strongly correlated electrons in two dimensional lattice that allows to consider the cuprates ($J \gg I$) and Sr_2RuO_4 ($J \ll I$) on the same footing. The singlet SC in the s -state is absent in the strong correlation limit, the triplet p -pairing occurs due to the ferromagnetic fluctuations and the singlet d -pairing is induced by the antiferromagnetic fluctuations. The reason why T_c in the cuprates is much higher than in Sr_2RuO_4 is the different gap anisotropies. For the p -state $\mathbf{k}$ -dependence of the gap results in the cancellation of the van-Hove singularity while for the d -state the

gap anisotropy permits large van-Hove singularity contribution in the equation for T_c . For the question what is so specific in the copper oxides for high- T_c superconductivity the possible answer may be as follows: it is the planar $Cu-O$ σ -bonding resulting in the strong antiferromagnetic $Cu-Cu$ interaction, that induced the singlet pairing with $d_{x^2-y^2}$ symmetry.

We thank N.M.Plakida for useful discussions and I.O.Baklanov for numerical calculations. The work was supported by the Russian Federal Program "Integration of high school education and science", grant N69.

*Present address: L.V.Kirensky Institute of Physics, Krasnoyarsk, 660036, Russia.

Electronic address: sgo@post.krascience.rssi.ru

- - - - -

- [1] Y.Maeno, H.Hasimoto, K.Yoshida, S.Nisshizaki, T.Fujita, F.Lichtenberg, *Nature* **372**, 532 (1994).
- [2] D.Pines, *Physica B* **163**, 78 (1990).
- [3] T.Oguchi, *Phys. Rev. B* **51**, 1385 (1995).
- [4] N.E.Bickers, D.J.Scalapino, R.T.Scaletar, *Int. J. Mod. Phys. B* **1**, 687 (1987).
- [5] T.M.Rice, H.Sigrist, *J. Phys. Cond. Matter* **7**, L 643 (1995).
- [6] I.I.Mazin, D.J.Singh, *Phys. Rev. Lett.* **79**, 733 (1997).
- [7] J.B.Goodenough, *Magnetism and chemical bond* (John Wiley and Sons, N.Y.-London, 1963).
- [8] E.Dagotto, *Rev. Mod. Phys.* **66**, 763 (1994).
- [9] A.P.Mackenzie, S.R.Julian, A.J.Diver *et al*, *Phys. Rev. Lett.* **76**, 3786 (1996).
- [10] D.J.Singh, *Phys. Rev. B* **52**, 1358 (1995).
- [11] T.M.Riseman, P.G.Kealey, E.M.Forgan, A.P.Mackenzie, L.M.Galvin, A.W.Tyger, S.L.Lee, C.Ages, D.McK.Paul, C.M.Aegerter, R.Cubitt, Z.Q.Hao, T.Akima, Y.Maeno, *Nature (London)* **396**, 242 (1998).
- [12] S.V.Tyablikov, *Methods of quantum theory of magnetism* (2-nd edition, Moscow, Nauka, 1975).
- [13] R.O.Zaitsev, V.A.Ivanov, *Fiz. tverdogo tela* (Sov. Solid State Physics) **29**, 2554 (1987).
- [14] N.M.Plakida, V.Yu.Yushanhai, I.V.Stasyuk, *Physica C* **160**, 80 (1989).
- [15] Yu.A.Izyumov, M.I.Katsnelson, Yu.N.Skryabin, *Magnetism of itinerant electrons* (Moscow, Nauka, 1994).
